

Fast Scale for Dementia

Stage	Stage Name	Characteristic
1	Normal Aging	No deficits whatsoever
2	Possible Mild Cognitive Impairment	Subjective functional deficit
3	Mild Cognitive Impairment	Objective functional deficit interferes with a person's most complex tasks
4	Mild Dementia	IADLs become affected, such as bill paying, cooking, cleaning, traveling
5	Moderate Dementia	Needs help selecting proper attire
6a	Moderately Severe Dementia	Needs help putting on clothes
6b	Moderately Severe Dementia	Needs help bathing
6c	Moderately Severe Dementia	Needs help toileting
6d	Moderately Severe Dementia	Urinary incontinence
6e	Moderately Severe Dementia	Fecal incontinence
7a	Severe Dementia	Speaks 5-6 words during day
7b	Severe Dementia	Speaks only 1 word clearly
7c	Severe Dementia	Can no longer walk
7d	Severe Dementia	Can no longer sit up
7e	Severe Dementia	Can no longer smile
7f	Severe Dementia	Can no longer hold up head

Fast Scale for Dementia

Stage	Stage Name	Characteristic
1	Normal Aging	No deficits whatsoever
2	Possible Mild Cognitive Impairment	Subjective functional deficit
3	Mild Cognitive Impairment	Objective functional deficit interferes with a person's most complex tasks
4	Mild Dementia	IADLs become affected, such as bill paying, cooking, cleaning, traveling
5	Moderate Dementia	Needs help selecting proper attire
6a	Moderately Severe Dementia	Needs help putting on clothes
6b	Moderately Severe Dementia	Needs help bathing
6c	Moderately Severe Dementia	Needs help toileting
6d	Moderately Severe Dementia	Urinary incontinence
6e	Moderately Severe Dementia	Fecal incontinence
7a	Severe Dementia	Speaks 5-6 words during day
7b	Severe Dementia	Speaks only 1 word clearly
7c	Severe Dementia	Can no longer walk
7d	Severe Dementia	Can no longer sit up
7e	Severe Dementia	Can no longer smile
7f	Severe Dementia	Can no longer hold up head

Palliative Performance Scale (PPSV2)

version 2

PPS Level	Ambulation	Activity & Evidence of Disease	Self-Care	Intake	Conscious Level
100%	Full	Normal activity & work No evidence of disease	Full	Normal	Full
90%	Full	Normal activity & work Some evidence of disease	Full	Normal	Full
80%	Full	Normal activity with Effort Some evidence of disease	Full	Normal or reduced	Full
70%	Reduced	Unable Normal Job/Work Significant disease	Full	Normal or reduced	Full
60%	Reduced	Unable hobby/house work Significant disease	Occasional assistance necessary	Normal or reduced	Full or Confusion
50%	Mainly Still/Lie	Unable to do any work Extensive disease	Considerable assistance required	Normal or reduced	Full or Confusion
40%	Mainly in Bed	Unable to do most activity Extensive disease	Mainly assistance	Normal or reduced	Full or Drowsy +/- Confusion
30%	Totally Bed Bound	Unable to do any activity Extensive disease	Total Care	Normal or reduced	Full or Drowsy +/- Confusion
20%	Totally Bed Bound	Unable to do any activity Extensive disease	Total Care	Minimal to sips	Full or Drowsy +/- Confusion
10%	Totally Bed Bound	Unable to do any activity Extensive disease	Total Care	Mouth care only	Drowsy or Coma +/- Confusion
0%	Death	-	-	-	-

Palliative Performance Scale (PPSV2)

version 2

PPS Level	Ambulation	Activity & Evidence of Disease	Self-Care	Intake	Conscious Level
100%	Full	Normal activity & work No evidence of disease	Full	Normal	Full
90%	Full	Normal activity & work Some evidence of disease	Full	Normal	Full
80%	Full	Normal activity with Effort Some evidence of disease	Full	Normal or reduced	Full
70%	Reduced	Unable Normal Job/Work Significant disease	Full	Normal or reduced	Full
60%	Reduced	Unable hobby/house work Significant disease	Occasional assistance necessary	Normal or reduced	Full or Confusion
50%	Mainly Still/Lie	Unable to do any work Extensive disease	Considerable assistance required	Normal or reduced	Full or Confusion
40%	Mainly in Bed	Unable to do most activity Extensive disease	Mainly assistance	Normal or reduced	Full or Drowsy +/- Confusion
30%	Totally Bed Bound	Unable to do any activity Extensive disease	Total Care	Normal or reduced	Full or Drowsy +/- Confusion
20%	Totally Bed Bound	Unable to do any activity Extensive disease	Total Care	Minimal to sips	Full or Drowsy +/- Confusion
10%	Totally Bed Bound	Unable to do any activity Extensive disease	Total Care	Mouth care only	Drowsy or Coma +/- Confusion
0%	Death	-	-	-	-

Palliative Performance Scale (PPSv2)					
PPS Level	Ambulation	Activity & Evidence of Disease	Self-Care	Intake	Conscious Level
100%	Full	Normal activity & work No evidence of disease	Full	Normal	Full
90%	Full	Normal activity & work Some evidence of disease	Full	Normal	Full
80%	Full	Normal activity <i>with</i> effort Some evidence of disease	Full	Normal or reduced	Full
70%	Reduced	Unable normal job/work Significant disease	Full	Normal or reduced	Full
60%	Reduced	Unable hobby/house work Significant disease	Occasional assistance necessary	Normal or reduced	Full or Confusion
50%	Mainly Sit/Lie	Unable to do any work Extensive disease	Considerable assistance required	Normal or reduced	Full or Confusion
40%	Mainly in Bed	Unable to do most activity Extensive disease	Mainly assistance	Normal or reduced	Full or Drowsy +/- Confusion
30%	Totally Bed Bound	Unable to do any activity Extensive disease	Total Care	Normal or reduced	Full or Drowsy +/- Confusion
20%	Totally Bed Bound	Unable to do any activity Extensive disease	Total Care	Minimal to sips	Full or Drowsy +/- Confusion
10%	Totally Bed Bound	Unable to do any activity Extensive disease	Total Care	Mouth care only	Drowsy or Coma +/- Confusion
0%	Death	-	-	-	-

Palliative Performance Scale (PPSv2)
version 2

PPS Level	Ambulation	Activity & Evidence of Disease	Self-Care	Intake	Conscious Level
100%	Full	Normal activity & work No evidence of disease	Full	Normal	Full
90%	Full	Normal activity & work Some evidence of disease	Full	Normal	Full
80%	Full	Normal activity <i>with Effort</i> Some evidence of disease	Full	Normal or reduced	Full
70%	Reduced	Unable Normal Job/Work Significant disease	Full	Normal or reduced	Full
60%	Reduced	Unable hobby/house work Significant disease	Occasional assistance necessary	Normal or reduced	Full or Confusion
50%	Mainly Sit/Lie	Unable to do any work Extensive disease	Considerable assistance required	Normal or reduced	Full or Confusion
40%	Mainly in Bed	Unable to do most activity Extensive disease	Mainly assistance	Normal or reduced	Full or Drowsy +/- Confusion
30%	Totally Bed Bound	Unable to do any activity Extensive disease	Total Care	Normal or reduced	Full or Drowsy +/- Confusion
20%	Totally Bed Bound	Unable to do any activity Extensive disease	Total Care	Minimal to sips	Full or Drowsy +/- Confusion
10%	Totally Bed Bound	Unable to do any activity Extensive disease	Total Care	Mouth care only	Drowsy or Coma +/- Confusion
0%	Death	-	-	-	-